

Recent Advances in the LEWICE Icing Model

William Wright, Vantage Group, LLC

Gene Addy, NASA Glenn

Peter Struk, NASA Glenn


Tadas Bartkus, Ohio Aerospace Institute

Glenn Research Center

Icing Branch

Page 1

at Lewis Field



CD-10-83242

Outline

- **Thermal Model Modifications**
- **Thermal Model Validation**
 - Comparison with Thermal Scaling Data
- **Conclusions from Thermal Analysis**
- **Mixed Phase Additions**
- **Mixed Phase Calibration**
 - Comparison with RatFac Data
- **Conclusions From Ice Crystal Analysis**



Thermal Model Additions

- **Myers Water Film Model**

$$\frac{\partial h}{\partial t} + \frac{\partial}{\partial x} \left[\frac{h^3}{3\mu_w} \left(\sigma \frac{\partial^3 h}{\partial x^3} + G_3 \frac{\partial h}{\partial x} - G_1 \right) + \tau_w \frac{h^2}{2\mu_w} \right] = \frac{\rho_a}{\rho_w} \beta V_\infty$$

- **Surface Water Shedding Model (calibrated)**

$$\frac{\dot{m}_{shed}}{\dot{m}_{runback,in}} = \frac{We - We_c}{We} \quad We = \frac{\rho_a V_a^2 x_k}{\sigma} \quad We_c = 200 + 5 * 10^5 x_k$$

- **Enhanced Evaporation**

- Chilton-Colburn analogy underestimates evaporation rate by 30%



Process for Comparison

- **Determine Internal Heat Transfer Coefficient from Dry Cases**
 - **All Cases Use Same Coefficients**

$$Nu = 0.004 \text{Re} \text{Pr}^{1/3} \left(\frac{z_n}{d_h} \right)^{-0.22} \left(\frac{x}{d_h} \right)^{-0.38}$$

- **External Heat Transfer Coefficient is Forced Laminar Where There is No Ice**
- **Run All Dry Cases To Ensure Correlation Matches**
- **Run Wet Cases for Validation**




Conditions Used For Thermal Comparison

Case	P(Pa)	V(m/s)	T(K)	LWC	MVD	AOA	t(min)
Warm Hold(Ref)	57295	92.7	264.5	0.5	20	0	7
Warm Hold(Scale)	98525	54.3	266.9	0.85	27.8	0	7
Descent(Ref)	69981	92.7	253.1	0.15	20	0	7
Descent(Scale)	97422	66.8	254.9	0.21	24.5	0	7
Cold Hold(Ref)	57295	92.7	247.4	0.15	20	0	10
Cold Hold(Scale)	98318	54.6	245.5	0.25	27.8	0	10



Warm Hold (Ref) - Dry



Glenn Research Center


Icing Branch

Page 6

at Lewis Field



Warm Hold (ref) - Wet



Glenn Research Center


Icing Branch

Page 7


at Lewis Field



Warm Hold (Ref) Ice Shape Comparison



Warm Hold (Re Scale) - Dry



Glenn Research Center


Icing Branch

Page 9

at Lewis Field



Warm Hold (Re Scale) - Wet



Glenn Research Center

Icing Branch

Page 10

at Lewis Field




Warm Hold (Re Scale) Ice Shape

- No Ice from Experiment nor from LEWICE



Descent (Ref)



Glenn Research Center


Icing Branch

Page 12


at Lewis Field



Descent (Ref)



Descent (Re Scale)



Glenn Research Center


Icing Branch

Page 14


at Lewis Field



Descent (Re Scale)



Cold Hold (Ref) - wet



Glenn Research Center


Icing Branch

Page 16


at Lewis Field



Cold Hold (Ref)



Cold Hold (Re Scale) - wet



Glenn Research Center


Icing Branch

Page 18

at Lewis Field



Cold Hold (Re Scale)



Observations from Thermal Analysis

- **Temperature Prediction is Very Good to Excellent for Most Cases**
- **Warm Hold Cases Show Predicted Runback Ice Forward of Experiment**
 - Peak Ice Thickness Higher for LEWICE
- **Descent and Cold Show Predicted Runback Ice Forms Slightly Behind Experiment**
 - Peak Thickness Higher for LEWICE, Especially Upper Surface
- **Ice in Experiment Grows Toward Leading Edge While LEWICE always grows Ice Normal to Surface**
- **Further Refinement of Runback Model May Be Necessary**
- **External Heat Transfer Coefficients for Residual Ice Shapes Need to Be Separately Validated**



Ice Breakup Model

- Breakup Threshold (Hauk)

$$V_{imp} \geq \frac{0.45}{\sin \alpha \sqrt{d}}$$

- Sticking Efficiency (Currie)

$$\frac{m_b}{m_o} = \left(1 - \xi \cos(\alpha_{imp})\right) \left(0.57 + 7.5 * 10^{-4} \left[V_{imp} \cos(\alpha_{imp})\right]^{1.5}\right)$$

$$\xi = -0.1425 + 47.292TWC - 1979.167TWC^2$$

– For $TWC < 0.12 \text{ kg/m}^3$ and $\xi = 0.14$ for $TWC > 0.12$



Conditions for Ice Crystal Comparison

Airfoil	Scan#	P	V	T	Sh	LWC	IWC	AOA	t(m)
Wedge	889	6.5	87.4	12.7	8.3	1.4	4.4	-6	3
Wedge	996	10	83.9	4.3	5.6	1.3	6.9	-6	3.5
Wedge	1003	10	84.1	3.8	5.2	1.9	7.3	-6	3.5
NACA 0012	796	6.5	86.2	7.2	5.9	0.6	4.9	0	3

Glenn Research Center


Icing Branch

Page 22


at Lewis Field




Ice Shape Prediction for Scan 996



Ice Shape Prediction for Scan 1003



Sticking Efficiency on Wedge at Various Particle Sizes



Glenn Research Center


Icing Branch

Page 25

at Lewis Field



Ice Thickness Prediction for Scan 796 (NACA0012)



Observations from Ice Crystal Comparison

- **Peak Thickness is Over Predicted by LEWICE while Extent is Under Predicted**
 - Additional Erosion Effects may be Needed
 - Improved Model for Reimpingement of Ice Crystals
- **Additional Data is Needed to Complete Model**

